

# BREAST RECONSTRUCTION REVISION

## Post-Surgical Guide



### INCISIONS

- Your incisions are closed in multiple layers with absorbable sutures
- The surface is sealed with a medical glue
- If you had an abdominal scar revision:
  - Your abdominal closure may feel tight. Keep your body flexed when in bed or relaxing in a recliner by placing two pillows behind your back and a pillow under your knees. When walking, you may need to bend at the knees and flex the hips to relieve tension the first week.

### DRAINS

- Use proper hand hygiene; such as a gel sanitizer on hands before caring for drain.
- Family member/friend strip/drain the JP bulbs and record accurate output using clear cups.
- Discard drainage in toilet only.
- Call Doctor for any increase in bright bloody drainage, cloudy drainage, increase in pain, loss of suction (non-depressed bulb), leaking around JP insertion site, or clogged bulb.
- If there is accidental dislodgement of drain: apply pressure with gauze and call Doctor **immediately**.
- Support drains with the Velcro belt provided by hospital.
- Drains will stay in until they are outputting about 20 ml or less per 24 hours.

### EMERGENCIES

\*\* After hours, call Medlink Paging Service at (512) 323-5465

- Any signs of infection, call Doctor IMMEDIATELY
  - Temperature greater than 101.00 degrees
  - Significant swelling
  - Tender, red, swollen, warm areas, pus, yellow or green drainage
  - Swollen/painful or darkened/blackened skin
- ABSOLUTELY NO ICE or HEATING pads to incisions/breasts

### SHOWER

- Shower daily with mild soap like Dove.
- Use a very soft washcloth, do not rub the area, pat area dry.

**Dr. Christine Fisher**  
1015 E 32<sup>nd</sup> St  
Plaza St. David, Suite 306  
Austin, TX 78705

Phone: (512) 815-0123  
Fax: (512) 861-6206  
After hours: (512) 323-5465

# BREAST RECONSTRUCTION REVISION

## Post-Surgical Guide

### ACTIVITY

- No strenuous activity for the week after surgery.
- Walk a lot. Stairs are okay.
- Sleep on back with bra for 4 weeks.

### MEDICATIONS

- Take two Aleve in the morning and two in the evening
- For additional pain control, take prescribed narcotic pain tablets every 4-6 hours as needed.
  - Prevent constipation by taking one dose of Miralax daily

**Dr. Christine Fisher**  
1015 E 32<sup>nd</sup> St  
Plaza St. David, Suite 306  
Austin, TX 78705

Phone: (512) 815-0123  
Fax: (512) 861-6206  
After hours: (512) 323-5465