

LIPOSUCTION

Post-Surgical Guide



LIPOSUCTION SITES

- Moderate bruising and swelling is normal at the sites from which the fat was removed.
- Moderate discomfort is normal and may last 1-2 weeks.
- If a large volume of fat was removed, we recommend wearing a compression garment or wrapping in an ACE bandage for 2 weeks after surgery.

EMERGENCIES

** After hours, call Medlink Paging Service at (512) 323-5465

- Any signs of infection, call Doctor IMMEDIATELY
 - Temperature greater than 101.00 degrees
 - Significant swelling
 - Tender, red, swollen, warm areas, pus, yellow or green drainage
 - Swollen/painful or darkened/blackened skin
- ABSOLUTELY NO ICE or HEATING pads to injection or harvest site.

SHOWER

- Shower daily with mild soap like Dove.
- Use a very soft washcloth, do not rub the area, pat area dry.

ACTIVITY

- Avoid strenuous activity for the first week after surgery.

MEDICATIONS

- Take two Aleve in the morning and two in the evening
- For additional pain control, take prescribed narcotic pain tablets every 4-6 hours as needed.
 - Prevent constipation by taking one dose of Miralax daily until function returns to normal.

Dr. Christine Fisher
1015 E 32nd St
Plaza St. David, Suite 306
Austin, TX 78705

Phone: (512) 815-0123
Fax: (512) 861-6206
After hours: (512) 323-5465