

# BREAST AUGMENTATION

## Post-Surgical Guide



### INCISIONS

- Your incisions are closed in multiple layers with absorbable sutures
- The surface is sealed with a medical glue
- Keep a clean, dry dressing over the incisions. Some drainage should be expected.

### EMERGENCIES

\*\* After hours, call Medlink Paging Service at (512) 323-5465

- Any signs of infection call Doctor IMMEDIATELY
  - Temperature greater than 101.00 degrees
  - Significant swelling
  - Tender, red, swollen, warm areas, pus, yellow or green drainage
  - Swollen/painful or darkened/blackened skin
- ABSOLUTELY NO ICE or HEATING pads to incisions/breasts

### BATHING

- Shower daily with a mild soap like Dove starting 24 hours after surgery.
- Use a very soft washcloth, do not rub the area, pat area dry.
- Do not soak breasts in bath or swim for 4 weeks post-surgery.
- In the rare case that you have drains:
  - Strip drains and record output twice per day and bring your log to your follow-ups.
  - Do not shower until the drains have been out for 48 hours; sponge bathe only until then.

### ACTIVITY

- Activity for the first 4-6 weeks after surgery:
  - Relax, rest, light activity within the home only for first week
  - No twisting, no lifting greater than 5-10 lbs for 4-6 weeks
  - No pressure on breasts
  - No side lying while sleeping for 6 weeks. Wear surgical bra while sleeping for 6 weeks.
  - May elevate arms to shower, but not above shoulders for first 2 weeks.
  - Avoid sitting, particularly for prolonged periods during the first 2 weeks.
  - Wear surgical bra 24/7 for 6 weeks.
  - You may be instructed in implant massage 2-3 weeks post surgery.

### MEDICATIONS

- Take two Aleve in the morning and two in the evening.

**Dr. Christine Fisher**  
2905 San Gabriel St. Suite 100  
Austin, TX 78705

**Phone: (512) 815-0123**  
Fax: (512) 861-6206  
After hours: (512) 323-5465

# BREAST AUGMENTATION

## Post-Surgical Guide

- For additional pain control, take prescribed narcotic pain tablets every 4-6 hours as needed.
  - Prevent constipation by taking one dose of Miralax daily.
- Take Singulair 10mg daily for 3 months to prevent capsular contracture.
- Complete antibiotic course started after surgery.

**Dr. Christine Fisher**  
2905 San Gabriel St. Suite 100  
Austin, TX 78705

**Phone: (512) 815-0123**  
Fax: (512) 861-6206  
After hours: (512) 323-5465