

MASTOPEXY (“BREAST LIFT”) / BREAST REDUCTION / ONCOPLASTIC BREAST RECONSTRUCTION

Post-Surgical Guide



INCISIONS

- Your incisions are closed in multiple layers with absorbable sutures
- The surface is may be sealed with a medical glue

DRAINS

- Use proper hand hygiene; such as a gel sanitizer on hands before caring for drain.
- Family member/friend strip/drain the JP bulbs and record accurate output using clear cups.
- Discard drainage in toilet only.
- Call Doctor for any increase in bright bloody drainage, cloudy drainage, increase in pain, loss of suction (non-depressed bulb), leaking around JP insertion site, or clogged bulb.
- If there is accidental dislodgement of drain: apply pressure with gauze and call Doctor **immediately**.
- Support drains with the Velcro belt provided by hospital.
- Your drains will stay in until they are outputting 20 ml per 24 hours or less. Expect to have the drains in about 1 week.

EMERGENCIES

** After hours, call Medlink Paging Service at (512) 323-5465

- Any signs of infection, call Doctor IMMEDIATELY
 - Temperature greater than 101.00 degrees
 - Significant swelling
 - Tender, red, swollen, warm areas, pus, yellow or green drainage
 - Swollen/painful or darkened/blackened skin
- ABSOLUTELY NO ICE or HEATING pads to incisions/breasts

SHOWER

- While drains are in place, sponge bathe with mild soap like Dove - DO NOT SHOWER.
- After drains are removed, shower daily with mild soap like Dove.
- Use a very soft washcloth, do not rub the area, pat area dry.

Dr. Christine Fisher
2905 San Gabriel St. Suite 100
Austin, TX 78705

Phone: (512) 815-0123
Fax: (512) 861-6206
After hours: (512) 323-5465

MASTOPEXY (“BREAST LIFT”) / BREAST REDUCTION / ONCOPLASTIC BREAST RECONSTRUCTION

Post-Surgical Guide

ACTIVITY

- Activity for the first 4-6 weeks after surgery:
 - No heavy lifting or strenuous activity for 2 weeks
 - No twisting, no lifting greater than 5-10 lbs for 4-6 weeks
 - No pressure on breasts
 - No side lying while sleeping for 6 weeks
 - May elevate arms to shower, but not above shoulders
 - Wear surgical bra for 6 weeks post-surgery. **No underwire bras** for 6 weeks.
 - May drive after 1 week if off all pain medications

MEDICATIONS

- Take two Aleve in the morning and two in the evening
- For additional pain control, take prescribed narcotic pain tablets every 4-6 hours as needed.
 - Prevent constipation by taking one dose of Miralax daily